(12) STANDARD PATENT (19) AUSTRALIAN PATENT OFFICE Title (54)prepartion thereof International Patent Classification(s) (51)CO7K 16/06 (2006.01) CO7K 16/18 (2006.01)

(11) Application No. AU 2002237586 82

Document FP3 Appl. No. 10/690,639

Pharmaceutical composition of F(AB)2 fragments of antibodies and method for the

C07K 16/24 (2006.01)

Data of Filing: 2002.02.28 (21)Application No: 2002237586

WIPO No: WO02/068475 (87)

Priority Data (30)

(74)

(33) Country Number (31)(32)Date 2001.03.05 09/798,076

Publication Date: 2002.09.12 (43)Publication Journal Date: 2003.03.06 (43)Accepted Journal Date: 2006.12.21 (44)

Applicant(s) (71)Juan Lopez De Silanes; Rita Mancilla Nava; Jorge Paniagua Solis

(72)Inventor(s) Paniagua Solis, Jorge; Mancilla Nava, Rita; Lopez De Silanes, Juan

Agent / Attorney Cullen & Co, Level 26 239 George Street, Brisbane, QLD, 4000



LETTERS PATENT

STANDARD PATENT

2002237586

I, Fatima Beattie, the Commissioner of Patents, grant a Standard Patent with the following particulars:

Name and Address of Patentee(s):

Juan Lopez De Silanes

Lucerna No. 7, Col., Jualez Delegacion Cuauhtemoc, D.F. 06600, Mexico

Pita Mancilla Nava

Calzada de Tlalpan No. 4687, Colonia Toriello Guerra. Delegacion Tlalpan, D.F.: 14950, Mexico

Jorge Paniagua Solis

Amores No. 1304, Colonia del Valle, Delegación Benifo Juarez, D.F. 03100, Mexico.

Name of Actual Inventor(s):

Paniagua Solis, Jorge; Mancilla Nava, Rita and Lopez De Silanes, Juan.

Title of Invention:

Pharmaceutical composition of F(AB)2 fragments of antibodies and method for the preparation thereof

Term of Letters Patent:

Twenty years from 28 February 2002

Priority Details:

Number

Date

Filed with

09/798,076

5 March 2001

US



Dated this 5th day of April 2007

Fatima Beattie

Commissioner of Patents

PATENTS ACT 1990

10

15

The claims defining the invention are as follows:

- 1. A method for preparing a composition of F(ab')₂ antibody fragments that is substantially free of whole antibodies, the method comprising the steps of:
- (a) generating a source of antibodies by immunising animals with a complex antigenic mixture of animal venom, extracting blood plasma that is the source of antibodies and contacting with pepsin at a pH of 3.2 ± 0.2 under conditions to prepare an antibody digest containing $F(ab')_2$ fragments and being completely free of non-hydrolyzed antibodies;
- (b) treating said digested antibodies by two steps of ammonium sulphate precipitation, with at least one step at about 16% to about 22% weight by volume of ammonium sulphate for at least 30 minutes at a temperature of 55 °C \pm 4 °C, next cooling to approximately 8 °C \pm 4 °C for at least two hours;
 - (c) clarifying the solution by filtering at 12, 8, 4 or 0.22 μ;
- (d) subjecting the obtained solution to a second precipitation at about 32% to about 38% weight by volume of ammonium sulphate at about pH 6.8 ± 0.5 , next leaving the solution to settle for at least 12 hours in refrigeration;
- (e) centrifuging the resulting suspension to eliminate the supernatant and recover the paste of F(ab')₂ fragments precipitated in step (d);
- (f) subjecting the obtained paste of F(ab')₂ fragments to dialysis to eliminate salts and low weight components and
- 20 (g) passing the solution through a sterile filter
 - 2. The method of claim 1, wherein said venom is produced by a scorpion.
 - 3. The method of claim 1, wherein said venom is produced by a snake.
 - 4. The method of claim 1, wherein said venom is produced by a spider.
- 5. The method of claim 1 or claim 2, wherein the venom is selected from the group of scorpions of the taxonomic family Butidae and is selected one of the group consisting of: Centruroides elegans-elegans, Centruroides exilicanda, Centruroides infamatus, Centruroides limpidus-tecomanus, Centruroides limpidus, Centruroides suffusus.

- 6. The method of claim 1 or claim 3, wherein the venom is selected from one of the snake genera: Bothrops, Crotalus, Agkistrodon, Lachesis Sistrurus, or the species Micrurus nigrosinctus.
- 7. The method of claim 1 or claim 4, wherein the venom is from the spider species:

 Lactrodectus mactans.
 - 8. The method of any one of claims 1 to 7, wherein the obtained composition of F(ab')₂ fragments is free of albumin, free of whole antibodies and free of viruses and pyrogens.
 - 9. A composition of F(ab')₂ fragments having features a, b and c as follows:
 - (a) the F(ab')₂ fragments are produced by the method of any one of claims 1 to 8,
 - (b) the F(ab')₂ fragments are free of whole antibodies, free of albumin, free of viral particles, and free of pyrogens,
- (c) the F(ab')₂ fragments are polyclonal, and capable of neutralising all the antigenic determinants present in complex animal venoms utilized to generate the source of antibodies of step (a).
 - 10. The composition of claim 9 which includes a pharmaceutically acceptable carrier.

20

Date: 6 December 2006